ANNEX 3 Danger Signs of the Newborn

The figures below show the danger signs of the newborn. CHT Members should <u>immediately</u> refer the baby to the health facility if they identify any of the following danger signs.

	J. J	
Feeding less or not feeding	Convulsion	High or very low
 at all Not able to suck at the breast when the mother tries to put the baby to the breast several times over a few hours The baby was feeding well after bith but has stopped feeding well now. 	 The baby's arms and legs become stiff. The baby stops breathing and becomes blue. Recurring movement of a part of the body like twitching of the mouth or blinking of the eyes. 	 temperature Temperature is below 35.5 Celsius Temperature is above 37.5 Celsius
		HANH SO CONTRACTOR
Local Infection	No movement or less	Fast or difficulty in
 Most common infection occurs in umbilicus, skin, and eyes. Pus coming out of the umbilicus and the eyes, and skin boils with pus. The skin where the stump is attached to is red. 	 movement The baby only moves when stimulated, or doesn't move at all with stimulation 	 Breathing "Fast breathing": breathing rate is 60 per minute or more. "Chest indrawing": the lower chest wall goes in when the child breathes in, and the upper chest and abdomen move out.
Yellow Skin		
 The baby shows yellow skin on face within 24 hours after delivery The baby develops yellow palms and soles (more than 24 hours old). 		